Vitra 10 for Sub Cyclo





Glaucoma Treatment



SubLiminal® Cyclophotocoagulation


SubCyclo® is a new nondestructive laser procedure for the treatment of glaucoma.

Based on the transcleral cyclophotocoagulation principle, it relies on the 810 nm SubLiminal® stimulation of the ciliary body, producing the aqueous humour, and the uveoscleral pathway, secondary route of the aqueous outflow.

This procedure can be performed thanks to the combined use of the Quantel Medical Vitra™ 810 laser and the Quantel Medical SubCyclo® probe.




Tip of the probe in contact with the sclera and positioned over the identified treatment area. Probe perpendicular to the eveball.



Laser treatment is delivered by moving the tip of the probe in a continuous sliding motion:

- Superior quadrants: From 9:30 to 12:00 and from 12:00 to 2:30
- Inferior quadrants: From 3:30 to 6:00 and from 6:00 to 8:30 The 3 and 9 o'clock meridians must be avoided.

Vitra™ 810 - SubLiminal Technology



Composed of a train of extremely short microsecond pulses of which the duration and the intervals are fully adjustable, this subthreshold treatment mode allows a precise management of the thermal effect on the targeted tissues and preserves the structures of the ciliary body.

SubCyclo® Laser Probe



The SubCyclo® laser probe features a unique needle design allowing a perfect and accurate probe positioning during the treatment which can be enhanced thanks to the transillumination technique. (1)

The tip of the needle also features a glass sphere ensuring an easy and comfortable sweeping motion.

Image courtesy of Dr. Pascale Hamard



Glaucoma Treatment


• SubCyclo® Clinical Data

A new treatment for advanced and end-stage glaucoma

Primary angle glaucoma, primary angle closure glaucoma, pseudo-exfoliation, neovascular glaucoma, glaucoma associated with steroid uses, uveitis and after keratoplasty surgery (2,3)

An efficient treatment procedure leading to a reduction in the medication regimen (2,4)

Significant and persistent reduction in IOP at 12 months (2)



30% decrease from baseline and less than 21 mmHg

Mean number of medication (2)

- Before treatment: 3.4 ± 1.7

- After treatment: **2.9** ± 1.6

SubCyclo® preserves the structures of the ciliary body (no destructive effects) and improves the uveoscleral drainage




Image courtesy of Dr. Nassima Benhatchi

After treatment:

- No anatomical damage or visible lesion have been observed by UBM (2)
- UBM shows a thin space between sclera and ciliary body which may correspond to the presence of suprachoroidal fluid ⁽²⁾

SubCyclo® avoids the side effects of the thermal cyclophotocoagulation procedure

No serious side effects such as intraocular inflammation after treatment ⁽²⁾ No severe complications such as phthisis bulbi, persistent hypotony, or cystoid macular edema ⁽²⁾

A repeatable treatment (2,4)

A safe treatment potentially allowing for earlier laser interventions

Vitra™ 810: A versatile laser

One single laser probe: Two glaucoma treatments

Quantel Medical SubCyclo® laser probe is uniquely designed with a removable footplate allowing implementing both SubLiminal® and thermal treatment procedures.









One laser: Two clinical applications

Although designed for the SubCyclo® laser therapy, Vitra™ 810 can also be used to perform all the usual 810 nm glaucoma and retina treatments.

When used in Retina treatment mode, Vitra™ 810 can be connected to various delivery systems, such as slit lamp, LIO, operating microscope and laser probes.











TECHNICAL SPECIFICATIONS

VITRA 810 nm

Laser Source: Laser diode technology

Wavelength: 810 nm Power at tissue up to: 3W

Pulse duration: 0.01 s to continuous Aiming beam: 635 - 650 nm

Repeat interval: 0.1 - 0.2 - 0.3 - 0.5 - 0.7 s

Emission modes: Single, repeat, continuous, painting

SubLiminal Mode (duty cycle from 5% to 35%) 18 (H) x 19.5 (W) x 30 (D) cm

Size: 18 (H) x 19.5 (W) x 30 (D) cm 7.1" (H) x 7.7"(W) x 11.8"(D) Weight: 5.6 kg - 12.3 lbs

Cooling: by Peltier effect
Power requirements: 100 to 240 VAC, 250 VA, 50/60 Hz

DELIVERY SYSTEMS

Slit lamp adapters

- Photocoagulation (Spot size: 100-1000 µm)

- Transpupillary Thermotherapy - TTT (Spot size: 1000-4000 µm)

Slit lamp compatibility: Quantel Medical slit lamp, Haag Streit 900 BM & BQ

and CSO SL 9800.

Laser indirect ophthalmoscope

Keeler Vantage Plus (external laser adaptor)

Laser probes

Straight 20, 23, 25 G Curved 20 G Flexible curved 23, 25 G Steerable 23, 25 G SubCyclo probe

OR microscope adapters

Zeiss type Leica type

© 2024.Quantel Medical, VitraTM 810, SubLiminal® and SubCyclo® are trademarks of Quantel Medical, member of Lumibird Medical Division. All rights reserved

BIBLIOGRAPHY

- 1- Agrawal P, Martin KR. Ciliary body position variability in glaucoma patients assessed by scleral transillumination. Eye. 2008;22:1499–1503.
- 2- Benhatchi N, Bensmail D, Lachkar Y. Benefits of SubCyclo laser therapy guided by high frequency ultrasound biomicroscopy in patients with refractory glaucoma. J Glaucoma 2019 Jun;28(6):535 539.
- 3- Keilani C, Benhatchi N, Bensmail D, Abitbol O, Amara A, Bluwol E, Graber M, Lachkar Y. Comparative effectiveness and tolerance of subliminal subthreshold transscleral cyclophotocoagulation with a duty factor of 25% versus 31.3% for advanced glaucoma. J Glaucoma. 2020 Feb;29(2):97-103.
- 4- Lutic I, Dragne C, Filip M, Filip A, Nicolae M, Moisescu R, Ungureanu I, Triantafyllidis G, Antonescu C. Subcyclo laser procedure results in patients with glaucoma. Rom J Ophthalmol. 2018 Oct-Dec;62(4):296-299.



XL_VIT810_CYCLO_BC6_AN_2
Artwork: www.overscan.fr







ManufacturerQuantel Medical

1 Rue du Bois Joli – CS40015 63808 Cournon d'Auvergne – FRANCE Tel.: +33 (0)4 73 745 745 Email: contact@lumibirdmedical.com ISO 13485 : 2016

Headquarters

Lumibird Medical 1 Rue du Bois Joli – CS40015 63808 Cournon d'Auvergne – FRANCE Tel.: +33 (0)4 73 745 745 QUANTEL MEDICAL
A brand of

